A Survey of the Pathogenic Fungus, *Batrachochytrium dendrobatidis*, at the Smithsonian Environmental Research Center, Anne Arundel County, Maryland.

Lauren D. Fuchs^{1†}, Todd A. Tupper^{2†*}, Tina Sephapur², Robert Aguilar³, Christine A. Bozarth²

¹George Mason University Department of Systems Biology 10900 University Blvd Manassas, VA 20110

²Northern Virginia Community College Department of Math, Science, and Engineering 5000 Dawes Avenue Alexandria, Virginia 22311

³Smithsonian Environmental Research Center Fish and Invertebrate Ecology Lab 647 Contees Wharf Road Edgewater, Maryland 21037

*Corresponding author: ttupper@nvcc.edu †Authors contributed equally to the development of the manuscript

Introduction

Batrachochytrium dendrobatidis (Bd) is a pathogenic species of chytrid fungus that produces flagellated, motile spores, which colonize keratinized epithelial cells in the skin of adult amphibians, and keratinized mouthparts of larval amphibians (Rachowicz and Vredenburg, 2004; Brutyn et al. 2012). It has been associated with declines and extirpations in over 200 amphibian species (Lips et al. 2006). Susceptibility to, and outcomes of *Bd* infection vary markedly both within and between species, ranging from no symptoms to mortality (Beebee and Griffiths 2005; Briggs et al. 2010). These inconsistencies are attributed to temperature (Berger et al. 2004), innate defenses (Harris et al. 2006; Woodhams et al. 2007), habitat (Kriger and Hero 2007; Rowley and Alford 2007), and host life history traits (Lips et al. 2003). Although Bd is found in a wide range of habitats and climates (Kilpatrick et al. 2010), catastrophic declines associated with Bd typically occur in cool, wet, and thermally consistent locations (Bielby et al. 2008; Murray et al. 2011; Berger et al. 2016). Such declines have been documented in Eastern Australia, Central America, and Europe (Berger et al. 1998; Lannoo et al. 2011; Johnson and Speare, 2003). In North America, observed declines have been restricted to the Western region of the United States (Longcore et al. 2007; Murray et al. 2009), with few *Bd*-related die-offs reported east of the Rocky Mountains (Petersen et al. 2016).

Studies in North America are generally sparse and opportunistic, showing inconsistencies in climatic determinants of *Bd* infectivity (Savage et al. 2011). Throughout the eastern United States, numerous studies (see Davidson and Chambers 2011; Huang and Wilson 2013; Lannoo et al. 2011; Longcore et al. 2007; Oullett et al. 2004; Peterson et al. 2016; Tupper et al. 2011) have identified the persistence of *Bd* in amphibians without associated population declines (Caruso and Lips 2013; Daszak et al. 2003; Muletz et al. 2014; Rothermel 2008). However, the mechanisms influencing

Catesbeiana 37(1): 32-40

the virulence of *Bd*, are poorly understood. A better understanding of its distribution and prevalence among amphibians is necessary to assess the potential threat of *Bd* on local amphibian populations (Daszak et al. 2003). To the best of our knowledge, *Bd* data from the Mid-Atlantic are limited; there are only two prior *Bd* studies conducted in Maryland (Grant et al. 2008; Hossack et al. 2010), and none have focused solely on anurans. Here we assessed the prevalence of *Bd* in anuran species at the Smithsonian Environmental Research Center (SERC), in Anne Arundel County, Maryland. Our primary objective was to assess the prevalence of *Bd* among anurans, thus enhancing existing, but limited information on *Bd* in Maryland. We also aimed to determine if *Bd* was confined to anurans of particular ecological guilds (Kriger and Hero 2007; Longcore et al. 2007; Tupper et al. 2011).

Methods

We opportunistically sampled for *Bd* at the Smithsonian Environmental Research Center (hereafter SERC; 38°53'17.41"N 76°33'15.52 W), in Anne Arundel County, Maryland (for more about SERC see Tupper et al. 2016) between 27 March and 27 June 2014 and 2015. We hand captured (while wearing new nitrile gloves) anurans and assaved skin surfaces via methods described by Hyatt et al. (2007) using sterile dry swabs (no. MW113, Medical Wire and Equipment Company, Durham, NC). We placed skin swabs in 1.5 mL microcentrifuge tubes, which were then frozen until molecular analyses. We followed the Purification of Total DNA from Animal Tissues protocol (Qiagen® DNeasy Blood & Tissue Kit, Valencia, CA) to extract DNA. To detect *Bd*, we prepared a 20 µL PCR reaction with 10 µL Sso AdvancedTM universal probes supermix (Bio-Rad, Hercules, CA), 200 nM each primer (ITS1-3Chytr and 5.8sChytr; Boyle et al. 2004), 250 nM MGB probe, and 2 µL of extracted DNA. We ran samples on a CFX96 Touch[™] Real-Time PCR Detection System (Bio-Rad, Hercules, CA) at 95°C for 3 min, followed by 45 cycles of 95°C for 30-sec and 55°C for 45-sec. We used positive and negative controls in DNA extraction and PCR amplification. Our goal was to determine the presence (not infection intensity) of Bd. Therefore, only one standard concentration was required as a positive control. We conducted qPCR analyses in triplicate on each sample to confirm a positive result. We considered a sample positive if it began to fluoresce before 40 cycles of our PCR reaction on at least two occasions. We used a Fisher's Exact test (Zar 2005) to assess differences in Bd detection rates between two broad ecological guilds of anurans: terrestrial/arboreal and aquatic (Table 1; Tupper et al. 2011) in minitab version 17 (ww.minitab.com). All nomenclature corresponds with Crother (2012).

Results

We opportunistically sampled 116 individuals across 11 species for *Bd*. Sixty-three (54.3 %) of the samples were collected in 2014 and 53 (45.7 %) were collected in 2015. Due to the opportunistic sampling scheme, sampling was not consistent between months and the monthly sampling distribution was not consistent between years. When pooled across years, five (4.3 %), 47 (40.5 %), 25 (21.6 %) and 39 (33.6 %) of the samples were collected in March, April, May and June, respectively. Seven of the 11 (63.6 %) species sampled tested positive for *Bd*, but only 12 of 116 (10.3 %) of the total samples contained *Bd*. Nine (75 %) of the positive detections were from 2014, and three (25 %) were collected in 2015. Six (50 %), three (25 %), and three (25 %) of the *Bd* positive samples were collected in May, June, and April, respectively. The largest proportion of detections (25%) occurred in the Pickerel Frog (*Lithobates palustris*) and Southern Leopard Frog (*Lithobates sphenocephalus*). The Eastern Cricket Frog (*Lithobates catesbeianus*; 16.7 %) and Green Frog (*Lithobates catesbeianus*; 10.5 %). We also identified *Bd* in Cope's Gray Treefrog (*Hyla chrysoscelis*) and American Toads (*Anaxyrus americanus*; < 1%). The Spring Peeper (*Pseudacris crucifer*), Eastern Spadefoot (*Scaphiopus holbrookii*), Wood Frog (*Lithobates sylvaticus*) and Gray Treefrog (*Hyla versicolor*) were

all *Bd* negative (Table 1). Significantly more detections of *Bd* occurred in the aquatic guild (16.9%; N = 59) than in the terrestrial and arboreal guild (0.04%; N = 57; Fisher's exact test P < 0.05).

Discussion

Although we detected *Bd* in 63.6 % of species sampled, the detection rate across individuals was low (10.3 %), and we did not observe visual signs of chytridiomycosis, the diseased state as a result of Bd infection (Muths et al. 2008). Our results are similar to studies conducted throughout the eastern United States, which have documented Bd in a wide range of amphibian species generally absent of clinical signs of chytridiomycosis (Grant et al. 2008; Pullen et al. 2010), and without associated catastrophic declines (Lannoo et al. 2011; Longcore et al. 2007; Petersen et al. 2016; Rothermel et al. 2008). Our work is also similar to other studies in that Bd was primarily detected in aquatic species but was not restricted to fully aquatic species (Longcore et al. 2007; Pearl et al. 2007; Rodriguez et al. 2009; Tupper et al. 2011). However, our results differ in terms of detection rates: we found one of the lowest detection rates across anuran species in the eastern United States (Davidson and Chambers 2011; Huang and Wilson 2013; Lannoo et al. 2011; Longcore et al. 2007; Petersen et al. 2016; Rothermel et al. 2008; Tupper et al. 2011; Table 2), the lowest in Maryland and, with the exception of Augustine and Neff (2016), the lowest detection rates when compared to similar studies conducted throughout Virginia. These comparisons include sites at the nearby Chesapeake and Ohio National Historic Park, Montgomery County, Maryland, and Huntley Meadows Park, Fairfax County, Virginia (Davidson and Chambers 2011; Grant et al. 2008; Goodman and Ararso 2012; Pullen et al. 2010; Hughey et al. 2014; Tupper et al. in progress). Results from Long Branch Nature Center, Arlington County, Virginia (Augustine and Neff 2016) must be interpreted cautiously, as the sample size was low (N = 25). In addition, sampling only spanned five genera and six species, with just three anuran species represented (N = 11).

Despite increased *Bd* sampling efforts in the eastern United States, certain areas still remain insufficiently surveyed (Olson et. al. 2013; GBDMP 2016). In addition to our study, only two other studies have investigated *Bd* in amphibians of Maryland (Grant et al. 2008; Hossack et al. 2010) and only Grant et al. (2008) sampled anurans. Grant et al. (2008) had a notably lower sample size (N = 53) than the present study, and only sampled three species (American Bullfrog, Pickerel Frog, and Green Frog) within a single genus (*Lithobates*). Therefore, our data add considerably to the understanding of *Bd* prevalence in Maryland across a wide range of anurans.

Although the presence of *Bd* throughout the eastern United States is well documented, it has yet to be associated with declines (Longcore et al. 2007; Grant et al. 2008; Pullen et al. 2010; Rothermel et al. 2008). Nevertheless, data implicate *Bd* as a primary source of certain dramatic amphibian declines worldwide (Berger et al. 1998; Collins and Crump 2009; Skerratt et al. 2007; Talley et al. 2015). The enigmatic nature of these declines stems from difficulties in demonstrating a causative link between disease and decline (Daszak et al. 2003). This difficulty is due to complex host-pathogen dynamics, and co-existing stressors (i.e. pollution, introduced species, increased UVB-radiation, and climate change) that may be implicated independently, or in conjunction with the pathogen (Beebee and Griffiths 2005). The extent to which each of these factors interacts with, and compounds the effects of *Bd* at the individual, community, and population level is not thoroughly understood, and hypotheses explaining these interactions are controversial (Blaustein et al. 2011). Our ability to interpret these complex interactions, and identify the potential threat of *Bd* on local amphibian populations relies heavily on continued surveillance, both at new and existing locations. We therefore recommend continued *Bd* monitoring using heightened biosecurity protocols (VHS 2016) at SERC and throughout the region.

Acknowledgements

Sara Ebrahimi and Coline Hay were essential to completion of fieldwork. We would like to also thank the following people for field assistance: Sonya Boldaji, Danny Boothe, David Graham, Kristin Jayd, and Lawanda Lumpkin. Lab work was completed at the Alexandria Campus of Northern Virginia Community College (Department of Math, Science, and Engineering). The Northern Virginia Community College Educational Foundation, the Smithsonian Environmental Research Center, and Dan Gustafson provided financial and logistical support. *Bd* sampling was approved by the Maryland Department of Natural Resources.

Table 1. Number and proportion of *Bd* positives by species at the Smithsonian Environmental Research Center. Superscript numbers following species are ecological guilds. 1 = terrestrial/arboreal, and 2 = aquatic. N = sample size.

Species	Ν	Bd Positive	% Positive
American Bullfrog ²	6	1	17
American Toad ¹	21	1	5
Cope's Gray Treefrog ¹	14	1	7
Eastern Cricket Frog ¹	18	4	22
Eastern Spadefoot ¹	8	0	0
Gray Treefrog ¹	1	0	0
Green Frog ²	19	2	11
Pickerel Frog ²	8	2	25
Southern Leopard			
Frog ²	4	1	25
Spring Peeper ¹	13	0	0
Wood Frog ²	4	0	0

Table 2. Proportion of *Bd* detections in anuran species sampled throughout Maryland and Virginia between 2008 and 2017. A range for % *Bd* Positive indicates that data were presented as percentages per species and site. * = Bedford, Campbell, Craig, Giles, Lynchburg City, Montgomery and Richmond City Counties. ** = Buckingham, Charles City, and Henrico Counties. N = Sample size.

Study	Location	Ν	% <i>Bd</i> Positive
Augustine & Neff 2016	Arlington County, VA	11	0
Davidson & Chambers 2011	Wise County, VA	41	14.6
Goodman & Ararso 2012	Prince Edward County, VA	103	7.8
Grant et al. 2008	Montgomery County, MD	53	17
Hughey et al. 2014	Throughout Western VA*	292	0-92
Present Study	Anne Arundel County, MD	116	10
Pullen et al. 2011	Throughout Central VA**	740	14.1

Literature Cited

- Augustine, L., and M. Neff. 2016. Determining the prevalence of amphibian chytrid fungus (*Batrachochytrium dendrobatidis*) and Ranavirus at Long Branch Nature Center in Arlington, Virginia. Catesbeiana 36: 35–42.
- Beebee, T. J. C., and R. A. Griffiths. 2005. The amphibian decline crisis: A watershed for conservation biology? Biological Conservation 125: 271–281.

Berger, L., R. Speare, P. Daszak, D. E. Green, A. A. Cunningham, C. L. Goggin, R. Slocombe, M. A. Ragan, A. D. Hyatt, K. R. McDonald, H. B. Hines, K. R. Lips, G. Marantelli, and H. Parkes. 1998. Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. Proceedings of the National Academy of Sciences of the United States of America 95, 9031–9036.

- Berger, L., R. Speare, H. B. Hines, G. Marantelli, A. D. Hyatt, K. R. McDonald, L. F. Skerratt, V. Olsen, J. M. Clarke, G. Gillespie, M. Mahony, N. Sheppard, C. Williams, and M. J. Tyler. 2004. Effect of season and temperature on mortality in amphibians due to chytridiomycosis. Australian Veterinary Journal 82: 31–36.
- Berger, L., A. A. Roberts, J. Voyles, J. E. Longcore, K. A. Murray, L. F. Skerratt. 2016. History and recent progress on chytridiomycosis in amphibians. Fungal Ecology 19: 89–99.
- Bielby, J., N. Cooper, A. A. Cunningham, T. W. J. Garner, and A. Purvis. 2008. Predicting susceptibility to future declines in the world's frogs. Conservation Letters 1: 82–90.
- Blaustein A. R., B. A. Han, R. A. Relyea, P. T. Johnson, J. C. Buck, S. S. Gervasi, and L. B. Kats. 2011. The complexity of amphibian population declines: understanding the role of cofactors in driving amphibian losses. Annals of the New York Academy of Sciences 1223: 108–119. doi: 10.1111/j.1749–6632.2010.05909.x.
- Boyle D.G., D.B. Boyle, V. Olsen, J.A.T. Morgan, and A.D. Hyatt. 2004. Rapid quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR assay. Diseases of Aquatic Organisms 60:141–148.
- Briggs, C. J., A. R. Knapp, and V. T. Vredenburg. 2010. Enzootic and epizootic dynamics of the chtryid fungal pathogen of amphibians. Proceedings of the National Academy of Sciences of the United States of America 107: 9695–9700. http://doi.org/10.1073/pnas.0912886107
- Brutyn, M., K. D'Herde, M. Dhaenens, P. Van Rooij, E. Verbrugghee, A. D. Hyatt, S. Croubels, D. Deforce, R. Ducatelle, F. Haesebrouck, A. Martel, and F. Pasmans. 2012. *Batrachochytrium dendrobatidis* zoospore secretions rapidly disturb intercellular junctions in frog skin. Fungal Genetics and Biology 49: 830–837.
- Caruso, N. M., and K. R. Lips. 2013. Truly enigmatic declines in terrestrial salamander populations in Great Smoky Mountains National Park. Diversity and Distributions 19: 38–38.
- Collins, J. P., and M. L. Crump. 2009. Extinction in our times: global amphibian declines. Oxford University Press, Oxford.

- Crother, B. I. 2012. Scientific and standard English names of amphibians and reptiles of North America North of Mexico, pp. 1–92, SSAR Herpetological Circular 39.
- Daszak, P., A. A. Cunningham, and A. D. Hyatt. 2003. Infectious disease and amphibian population declines. Diversity and Distributions 9: 141–150.
- Davidson, S. R. and D. L. Chambers. 2011. Occurrence of *Batrachochytrium dendrobatidis* in amphibians of Wise County, Virginia, USA. Herpetological Review 42: 214–216.
- GBDMP (Global *Bd*-Mapping Project- *Bd*-Maps). 2016. http://www.bdmaps.net/surveillance/s_country.asp. (Accessed 3 June 2016).
- Goodman, R.M., and Y.T Ararso. 2012. Survey of ranvirus and the fungus *Batrachochytrium denodrobatidis* in frogs of central Virginia, USA. Herpetological Review 43:78–80.
- Grant, E. H. C., L. L. Bailey, J. L. Ware, and K. L. Duncan. 2008. Prevalence of the amphibian pathogen *Batrachochytrium dendrobatidis* in stream and wetland populations in Maryland, USA. Applied Herpetology 5: 233–241.
- Harris, R. N., T. Y. James, A. Lauer, M. A. Simon, and A. Patel. 2006. Amphibian pathogen *Batrachochytrium dendrobatidis* is inhibited by the cutaneous bacteria of amphibian species. EcoHealth 3:53–56.
- Hossack, B. R., M. J. Adams, E. H. C. Grant, C. A. Pearl, J. B. Bettaso, W. J. Barichivich, W. H. Lowe, K. True, J. L. Ware, and P. S. Corn. 2010. Low prevalence of chytrid fungus (*Batrachochytrium denodrobatidis*) in amphibians of U.S. headwater streams. Journal of Herpetology 44: 253–260.
- Huang, R. and L. A. Wilson. 2013. *Batrachochytrium denodrobatidis* in amphibians of the Piedmont and Blue Ridge provinces in northern Georgia, USA. Herpetological Review 44: 95–98.
- Hughey, M.C., M.H. Becker, J.B. Walke, M.C. Swartwout, and L. K. Belden. 2014. *Batrachochytrium denodrobatidis* in Virginia amphibians within and among site variation in infection. Herpetological Review 45:428–438.
- Hyatt, A. D., D. G. Boyle, V. Olsen, D. B. Boyle, L. Berger, D. Obendorf, A. Dalton, K. Kriger, M. Hero, H. Hines, R. Phillott, R.Campbell, G. Marantelli, F. Gleason, and A. Colling. 2007. Diagnostic assays and sampling protocols for the detection of *Batrachochytrium dendrobatidis*. Diseases of Aquatic Organisms 73:175–192.
- Johnson, M. L. and R. Speare. 2003. Survival of *Batrachochytrium dendrobatidis* in water: quarantine and disease control implications. Emerging Infectious Diseases 9:922-925.
- Kilpatrick, A. M., C. J. Briggs, and P. Daszak. 2010. The Ecology and impact of chytridiomycosis: an emerging disease of amphibians. Trends in Ecology and Evolution 25: 109–118.

- Kriger, K. M., and J M. Hero. 2007. The chytrid fungus *Batrachochytrium dendrobatidis* is non-randomly distributed across amphibian breeding habitats. Divers Distrib. 13:781–788.
- Lannoo, M. J., C. Petersen, R. E. Lovich, P. Nanjappa, C. Phillips, J. C. Mitchell, and I. Macallister. 2011. Do frogs get their kicks on Route 66? Continental U.S. Transect reveals spatial and temporal patterns of *Batrachochytrium dendrobatidis* infection. PLoS ONE 6: e22211. doi: 10.1371/journal.pone.0022211
- Lips, K. R., J. D. Reeve, and L. R. Witters. 2003. Ecological traits predicting amphibian population declines in Central America. Conservation Biology 17:1078–1088.
- Lips, K. R., F. Brem, R. Brenes, J. D. Reeve, R. A. Alford, J. Voyles, C. Carey, L. Livo, A.P. Pessier, and J. P. Collins. 2006. Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. Proceedings of the National Academy of Sciences of the United States of America 103: 3165–3170.
- Longcore, J. R., J. E. Longcore, A. P. Pessier, and W. A. Halteman. 2007. Chytridiomycosis widespread in anurans of northeastern United States. Journal of Wildlife Management 71: 435–444.
- Muletz, C., N. M. Caruso, R. C. Fleischer, R. W. McDiarmid, and K. R. Lips. 2014. Unexpected rarity of the pathogen *Batrachochytrium dendrobatidis* in Appalachian *Plethodon* salamanders: 1957–2011. PLoS ONE 9(8): e103728. doi: 10.1371/journal.pone.0103728
- Murray, K. A., D. Rosauer, H. McCallum, and L. F. Skerratt. 2011. Integrating species traits with extrinsic threats: closing the gap between predicting and preventing species declines. Proceedings of the Royal Society B: Biological Sciences 278: 1515–1523.
- Muths, E., D. S. Pilliod, and L. J. Livo. 2008. Distribution and environmental limitations of an amphibian pathogen in the Rocky Mountains, USA. Biological Conservation 141: 1484–1492.
- Olson, D. H., D. M. Aanensen, K. L. Ronnenberg, C. I. Powell, S. F. Walker, J. Bielby, T. W. J. Gerner, G. Weaver, The Bd Mapping Group, and M. C. Fisher. 2013. Mapping the global emergence of *Batrachochytrium dendrobatidis*, the amphibian chytrid fungus. PLoS ONE 8(2): e56802. doi: 10.1371/journal.pone.0056802.
- Oullet, M., I. Mikaelian, B. D. Pauli, J. Rodriguez, and D. M. Green. 2005. Historical evidence of widespread chytrid infection in North American amphibian populations. Conservation Biology 19: 1431–1440.
- Pearl, C. A., E. L. Bull, D. E. Green, J. Bowerman, M. J. Adams, A. Hyatt, and W. H. Wente. 2007. Occurrence of the amphibian pathogen *Batrachochytrium dendrobatidis* in the Pacific Northwest. Journal of Herpetology. 41:145–149.
- Petersen, C. E., R. E. Lovich, C. A. Phillips, M. J. Dreslik, and M. J. Lannoo. 2016. Prevalence and seasonality of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* along widely separated longitudes across the United States. EcoHealth pp 1-15. DOI 10.1007/s10393-016-1101–4.

- Pullen, K. D., A. M. Best, and J. L. Ware. 2010. Amphibian pathogen *Batrachochytrium dendrobatidis* prevalence is correlated with season and not urbanization in central Virginia. Diseases of Aquatic Organisms 91: 9–16.
- Rachowicz, L. J. and V. T Vredenburg. 2004. Transmission of *Batrachochytrium dendrobatidis* within and between amphibian life stages. Diseases of aquatic organisms 61: 75-83.
- Rodriguez, E. M., T. Gamble, M. V. Hirt, and S. Cotner. 2009. Presence of *Batrachochytrium dendrobatidis* at the headwaters of the Mississippi River, Itasca State Park, Minnesota, USA. Herpetological Review 40:48–50.
- Rothermel, B. B., S. C. Walls, J. C. Mitchell, C. K. Dodd Jr., L. K. Irwin, D. E. Green, V. M. Vasquez, J. W. Petranka, and D. J. Stevenson. 2008. Widespread occurrence of the amphibian chytrid fungus *Batrachochytrium deondrobatidis* in the southeastern USA. Diseases of Aquatic Organisms 82: 3–18.
- Rowley, J. J. and R. A. Alford. 2007. Behaviour of Australian ranforest stream frogs may affects the transmission of chytridiomycosis. Diseases of Aquatic Organisms 77: 1–9.
- Savage, A. E., M. J. Sredl, and K. R. Zamudio. 2011. Disease dynamics vary spatially and temporally in a North American amphibian. Biological Conservation 144: 1910–1915.
- Skerratt, L.F., L. Berger, R. Speare, S. Cashins, K. R. McDonald, A.D. Phillott, H. B. Hines, and N. Kenyon. 2007. Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. EcoHealth 4: 125–134.
- Talley, B. L., C. R. Muletz, V. T. Vredenburg, R. C. Fleischer, and K. R. Lips. 2015. A century of *Batrachochytrium dendrobatidis* in Illinois amphibians (1888–1989). Biological Conservation 182: 254–261.
- Tupper, T.A. J.W. Streicher, S.E. Greenspan, B.C. Timm. 2011 and R.P Cook. Detection of *Batrachochytrium dendrobatidis* in anurans of Cape Cod National Seashore, Barnstable County, Massachusetts, USA. Herpetological Review 42: 62–65.
- Tupper, T. A., R. Aguilar, T. Sepahpur, C. Hay, and S. Ebrahimi. 2016. A survey of amphibians at the Smithsonian Environmental Research Center, Anne Arundel County, Maryland. Banisteria 46: 12–24.
- Tupper, T.A., L.D. Fuchs, C. Love, D. Lawlor and C.A. Bozarth. In progress. A survey of *Batrachochytrium dendrobatidis* at Huntley Meadows Park in Fairfax County, Virginia.
- VHS (Virginia Herpetological Society). 2016. Infectious diseases of our native herps and disinfection protocols. http://virginiaherpetologicalsociety.com/disease/index.htm. Accessed May 2016.
- Woodhams, D. C., K. Ardipradja, R. A. Alford, G. Marantelli, L. K. Reinert, and L. A. Rollins-Smith. 2007. Resistance to chytridiomycosis varies among amphibian species and is correlated with skin peptide defenses. Animal Conservation 10: 409–417.
- Zar, J. H. 2010. Biostatistical Analysis. 5th Edition. Prentice Hall, New York, NY. 960 pp.